Comparison of predator-parasitoid-prey interaction models for different host plant qualities

Carolina Sanchez¹, Manuel Gamez^{2*}, Francisco J. Burguillo³, József Garay⁴, Tomás Cabello¹

¹Department of Biology and Geology, University of Almeria, Almeria, Spain

²Department of Mathematics, University of Almeria, Almeria, Spain

³Department of Physical Chemistry, University of Salamanca, Salamanca, Spain

⁴Department of Plant Systematics, Ecology and Theoretical Biology, L. Eötvös University, Budapest, Hungary

* corresponding author e-mail: mgamez@ual.es

Keywords: phytophagous-parasitoid-predator system, Lotka-Volterra model, cultivar, *Nesidiocoris tenuis, Bemisia tabaci, Trichogramma achaeae*

Abstract: Population dynamics models suggest that the over-all level of resource productivity plays an important roles in community dynamics. One such factor of resource productivity the quality of the host plant, which can determine the effectiveness of entomophagous (predatory and parasitoids species) by altering the growth rate of the phytophagous population via effects on fecundity, survival, and rate of development. These effects have been studied in relation to the distribution of host plants and their physiological state. However, few studies have considered the differences among plant cultivars.

The objective of this study was to identify a continuous-time dynamic model, to describe the effects of different tomato cultivars on a one predator-two prey model. The experiment was carried out under greenhouse conditions using ten tomato cultivars, with the predatory species *Nesidiocoris tenuis* (Reuter) (Insecta, Hemiptera, Miridae) and two prey species: the phytophagous species *Bemisia tabaci* (Gennadius) (Insecta, Hemiptera, Aleyrodidae) and the parasitoid species *Trichogramma achaeae* Nagaraja & Nagarkatti (Insecta, Hymenoptera, Trichogrammatidae); the latter was used as the intraguild-prey.

Using the software SIMFIT we found that a three-dimensional Lotka-Volterra type system could be well fitted to the data, estimating the phytophagous species' growth rate, the parasitoid and predator mortality rates, the predation and parasitism rates, and the parasitoid emergence rate according to the cultivar type.

The results showed an important effect of the host plant quality,by cultivar, on intraguild predation, resulting in important changes in the dynamics of phytophagous populations. These results are also discussed in relation to their importance in the biological control of pest species in greenhouse crops.

Introduction

Biological control programmes in European greenhouse tomato crops depend on the use of *Nesidiocoris tenuis* (Reuter) or *Macrolophus pygmaeus* (Hem.: Miridae) to control the sweet potato whitefly, *Bemisia tabaci* (Gennadius) (Hem.: Aleyrodidae) (Castañe et al. 2007; Calvo et al. 2012) and on *Trichogramma achaeae* Nagaraja and Nagarkatti to control the South American tomato moth, *Tuta absoluta* (Meyrick) (Lep.: Gelechiidae) (Cabello et al. 2012, Oliveira et al. 2017).

N. tenuis is an omnivorous predator (Sánchez 2008), and *T. achaeae* is a parasitoid species that attacks lepidopteran eggs (Cabello et al. 2009, Polaszek et al. 2012). Thus, for tomato crops in greenhouses, an important intraguild predation (IGP), exists with *N. tenuis* as the IG-predator and immature stages of *T. achaeae*, inside the host eggs, as the IG-prey (Cabello et al. 2015). It should be noted that the use of different entomophagous species in greenhouse biological control programmes is relatively common (Vila and Cabello 2014). This can lead to direct or indirect interactions such as apparent competition, IGP and resource competition (Janssen et al., 1988). These interactions can have a global impact on the efficiency of biological control agents (Yano 2015, Messelink et al. 2013). Interest in these interactions has resulted in a significant amount of both theoretical and applied research. However, studies that show the impact of IGP in biological control are limited (Janssen et al. 1988, Rosenheim and Harmon 2006).

Evaluation of the efficiency of biological control agents, - in IGP conditions-, is difficult. Dynamic mathematical Lotka-Volterra type models have been proposed and used for this purpose (Holt and Polis 1997, Berryman 2004, Varga et al. 2010, Gallego et al. 2013).

The objective of this study was to find a continuous-time dynamic model, to describe the effects of different tomato cultivars on a one predator-two prey model.

Mathematical models applied to natural enemies have developed extensively over (the past 90 years), from primarily theoretical works, to more recent applied practices (Hawkins and Cornell, 2004).

This objective is being realized in collaboration with industrial producers of biological agents, since both the selection and use of the agent are relevant to applications by farmers and agronomists.

2. Material and Methods

2.1. Biological material

T. achaeae were reared at the Applied Entomology Laboratory at the University of Almeria (Spain) for 20 generations before the trial. This species was reared using the factitious host *Ephestia kuehniella* Zeller (Lep.: Pyralidae) in plastic containers (1 litre) following the methodology used by Cabello (1985). For that purpose, 12,500 *E. kuehniella* eggs (irradiated with ultraviolet light) were added to a piece of cardboard (13 x 10.5 cm) and placed in each container. Later, parasitoid adults at a ratio of 1:4 were introduced. The environmental conditions for the growth of the parasitoid were $25\pm1^{\circ}$ C, at 60-80% relative humidity (RH) with a photoperiod 16:9h (L:D). *N. tenuis* adults were

provided by Nesicontrol ® (Agrobio, La Mojonera, Almería, Spain). The *E. kuehniella* irradiated eggs were provided by Biotop (Valbonne, France). The eggs were refrigerated at 7°C to preserve them, until their use for both rearing the parasitoids and the greenhouse trial. Finally, *B. tabaci* adults were provided, by Koppert Biological Systems S.L. (Águilas, Murcia, Spain).

2.2. Experimental design

The experiment was designed random complete blocks with four repetitions and one factor (tomato cultivar) as the treatment (to 10 levels).

The trial was conducted from April to June 2015 in an experimental multitunnel greenhouse, with a 4,896 m² surface area in the TECNOVA Technology Centre (Viator, Almería, Spain). Four isolated modules (15 x 8 m) were built in the greenhouse using plastic film (1.0 x 0.5 mm). Each of these modules corresponded to one blocks in the trial design. Each module was divided into 10 plots (12m²). On April 7th , 18 of the following seedling cultivars were transplanted into each plots: Bronco®, and Pasadena® (Ramiro Arnedo S.A., Almería, Spain); Creativo®, and Genio® (Hm.Clause Ibérica S.A., Almería, Spain); Montenegro®, and Ramyle ® (Rijk Zwaan Ibérica S.A., Almería, Spain); Byelsa® (Semillas Fitó S.A., Almería, Spain); Caniles® (Zeraim Ibérica S.A., Almería, Spain); Rebelión® (Vilmorin Ibérica S.A., Almería, Spain) ; and Vernal® (Enza Zaden Investigation Centre S.L.U). Commercial crop management was used in the area, with an exception where no pesticide treatment was used.

All plots were artificially infested with the pest species, *B. tabaci*, on April 8th (1 day after transplanting, DAT) and April15th (8 DAT), at a dose of 30 adults/m². The natural enemies were released as follows: one release of *N. tenuis* on April 23rd (16 DAT) at a dose of 4 adults/m², and 4 releases of the parasitoid *T. achaeae* 241 parasitized eggs/m² every fifteen days, beginning on April 14th (17 DAT).

2.3. Sampling

Sampling was conducted every two weeks, a total of 5 times, beginning May 8th (31 DDT) for the pest species *B. tabaci* (nymph stage) and for the predator species *N. tenuis* (adult stage). For this purpose 4 plants were randomly chosen and the 10 last apical leaves of each plant were visually examined, for the number of pest species nymphs and predator adults. The 4 randomly chosen plants in each block and plot were used in subsequent samplings. The parasitoid *T. achaeae* was evaluated through the sentinel eggs method. To accomplish this, 4 cardboard pieces were placed in each block and plot, using the same plants sampled for *N. tenuis* and *B. tabaci*, to which we had added approximately 200 irradiated *E. kuehniella* eggs. This sampling was also conducted every two weeks, a total of 4 times, starting on May 7th (30 DAT). The eggs were exposed for 7 days in the greenhouse. They were later taken to the laboratory, and preserved at 25°C for an additional 7 days prior to quantification of eggs parasitized by *T. achaeae*. The phenological development of the plants during the trial period was also evaluated.

2.4.Data Analysis

The number of *B. tabaci* nymphs, *N. tenuis* adults, and parasitized eggs for each sampling event were initially expressed in square metre values. Later, those data were expressed as insect-day accumulated values (IDA). This index, proposed by Ruppel (1983), was applied to evaluate the total impact of pests at a given period of time. IDA

data were analysed with GLM univariate analysis for the cultivar factor. Analyses were conducted using IBM SPSS 23 statistical software (IBM 2014).

2.5. Mathematical model

Figure.- 1 shows the network interactions used in our model according to the nomenclature applied by Mill (2006). Its mathematical expression is shown below:

Pest species
$$x_1' = x_1 (m_1 - \alpha \cdot x_1 - \gamma_1 \cdot x_3)$$

Parasitoid species $x_2' = x_2 (m_2 - \gamma_2 \cdot x_3)$ (1)
Predator species $x_3' = x_3 (-m_3 + \overline{\gamma}_1 \cdot x_1 + \overline{\gamma}_2 \cdot x_2),$

where x_1 , x_2 , and x_3 are the densities (number / m²) of pest (phytophagous), parasitoid and predator species, respectively. According to the terminology of Abrams (2012), m_1 is the intrinsic growth rate of the pest (prey 1); m_2 is the increase rate of the parasitoid species (prey 2), due to the intrinsic growth rate and releasing rate; m_3 is the death rate of the predator in the absence of prey; α is the coefficient of intraspecific competition for pest species; γ_1 and γ_2 are the slopes of the predator's functional response when they kill the pest species (prey 1) and the parasitoid species (prey 2), respectively; and $\overline{\gamma}_1$ and $\overline{\gamma}_2$ are the slopes of the predator's numerical response when they kill and eat the pest species (prey 1) or the parasitoid species (prey 2), respectively.

Figure 1. Network of interactions considered in the model phytofagous – parasitoid – predator (P = pest *Bemisia tabaci*; T = parasitoid *Trichogramma achaeae* and N = predator *Nesidiocoris tenuis*); the linking arrows show benefits (+) and losses (–).



Using the statistical software SIMFIT version 2017 (Bardsley 2017), the system of equations (1) was fitted to the data corresponding to the number of *N. tenuis* adults , *B. tabaci* nymphs , and immature stages of *T. achaeae* per square metre in each sampling event for the three variables used: Byelsa, Caniles and Rebelion.

2.6. Optimal control of the system

Based on (1), we consider the control system

$$x' = F(x_1, x_2, x_{31}, u) = \begin{pmatrix} x_1(m_1 - \alpha \cdot x_1 - \gamma_1 \cdot x_3) \\ x_2(m_2 - \gamma_2 \cdot x_3) + u \\ x_3(-m_3 + \overline{\gamma}_1 \cdot x_1 + \overline{\gamma}_2 \cdot x_2) \end{pmatrix},$$

where control function u describes the time-dependent rate of release of parasitoids, realizing that the biological control of the pest T. *absoluta* is not considered in this model. Fix a time interval $[t_1, t_2]$, and for each $\varepsilon > 0$, let $U_{\varepsilon}[t_1, t_2]$ be the set of piece-wise constant, ε -small controls $0 \le u(t) \le \varepsilon$ ($t \in [t_1, t_2]$).

In order to keep the pest above a level *K*, minimizing the total release of agent, we have to solve the following optimal control problem:

$$\Psi(u) = \int_{t_1}^{t_2} c u(t) dt \to \min,$$

$$u \in U_{\varepsilon}[t_1, t_2],$$

$$\dot{x} = F(x_1, x_2, x_3, u_2, u_3), \quad x(0) = x^0,$$

$$g(x_1(t), x_2(t), x_3(t)) \ge 0 \quad (t \in [t_1, t_2]),$$
(2)

where function $g(x_1, x_2, x_3) = x_2 - K$ measures the level of the parasitoid, in comparison to a threshold value, below which the parasitoid .cannot control the pest *T. absoluta*. For the solution, the toolbox developed for MatLab by Banga *et al.* (2005) and Hirmajer *et al.* (2009) was applied.

3. Results

3.1. Effects of cultivars on populations of the three species

The cultivar was observed to have a significant effect on the N. tenuis predator's IDA for the pest species B. tabaci and the parasitoid species T. achaeae (F = 6.268, df = 9/150, P < 0.01; F = 8.589, df = 9/150, P < 0.01; F = 8.263, df = 9/150, P < 0.01, respectively) (Figure 2). In Figure 2A, all the predator IDA values were relatively high and didn't differ significantly, except for the cultivar Rebelion. The pest species had the lowest IDA values (Figure 2B) for the following three cultivars: Ramyle, Byelsa and Pasadena. Meanwhile, cultivar Rebelion showed the highest incidence of the pest species, and the remaining cultivars had intermediate IDA values. Finally, Figure 2C shows that the cultivar Byelsa exhibited the lowest incidence of the immature stages of the parasitoid, whose values differed significantly from those corresponding to the Bronco, Rebelion and Caniles cultivars.

Figure 2: Averages of insect-day accumulated (IDA) for *Nesidiocoris tenuis* adults (A), *Bemisia tabaci* nymphs (B), and immature stages of *Trichogramma achaeae* (C) in greenhouse tomato crops by cultivar (whiskers represent SE).



3.2. Population dynamic models

From the 10 cultivars previously evaluated, we selected Byelsa and Rebelion, which exhibited more extreme values of incidence regarding phytophagous species, predator, and parasitoid, and Caniles, a cultivar with an intermediate value.

Mathematical model (1) was well-fitted to the data for cultivars Byelsa and Caniles, but not to Rebelion, as shown in the corresponding curves in Figure 3. From this, we can conclude that the plants corresponding to the cultivar Rebelion presented a very low predator incidence, indicating that it had some impact on the pest species population, but not on the parasitoid population.

Caniles (Figure 3B) has very similar values to those of Byelsa and fits the model. However, the population of T. achaeae decreases due to intraguild predation before the value of B. tabaci (prey 1) declines

Finally, Rebelion could not be adjusted to our model. In this case, the predator population was very small, thus there was no effect on the *T. achaeae* population.

Figure 3: Densities obtained from the fitted model: prey species = *Bemisia tabaci* (nymph stage), parasitoid species = *Trichogramma achaeae*, and predatory species = *Nesidiocoris tenuis*, in greenhouse tomato crop for the cultivars: Byelsa (A), Caniles (B), and Rebelion (C).



Table 1: Fitting of the two-prey- one-predator model to the data of greenhouse tomato crop data for Byelsa and Caniles cultivars.

Cultivar	Fitting parameters (average ± SE)								Statistical parameters		
	<i>m</i> ₁	m ₂	<i>m</i> ₃	α	<i>γ</i> 1	<i>γ</i> 2	$\overline{\gamma}_1$	$\bar{\gamma}_2$	d.f.	r^2	Р
Byelsa	1.785	2.222	7 10 -4	0.224	1.318	1.811	0.131	2 10 -4	13	0.976	< 0.05
	(±0.166)	(±0.15)	(±0.256)	(±0.04)	(±0.214)	(±0.439)	(±0.011)	(±0.211)			
Caniles	1.605	3.584	7 10 -4	0.178	1.602	8.654	0.097	0.120	13	0.952	< 0.05
	(±0.140)	(±0.284)	(±0.164)	(±0.032)	(±0.293)	(±1.841)	(±0.011)	(±0.049)			

3.3. Solution of the optimal control problem

Based on the previous results, we conclude that the predator N. tenuis (adult stage) is necessary for the control of the pest species *B. tabaci* (nymph stage), but has adverse side effects on the immature stages of the parasitoid *T. achaeae* used for the control of a different pest species, *T. absoluta*, not considered in the model. From a theoretical point of view, we had the following question: How should the release rate of the parasitoid increase to maintain sufficient control of the pest *T. absoluta*? For that purpose, and based on previously published studies (e.g., Cabello et al. 2012, 2015), we determined that an appropriate density of parasitoids (immature stages) within crop plots was 100 individuals per square metre. Values lower than this density can cause damage due to the pest species *T. absoluta* (Figure 4A). The results (Figure 4B) show that, to maintain the aforementioned threshold value, it is necessary to carry out 7 additional releases in the crop cycle, for an overall density of 80 specimens/m2.

Figure 4: For $[t_1, t_2] = [4,10]$, the optimal trajectory (A) and control of the problem (B) for the two-prey-one-predator model using greenhouse tomato crop data for the Byelsa cultivar.



3.4. Stable coexistence of the population system. A theoretical point of view

As outlined in the previous section, for the considered species, an optimal release of the parasitoid can be obtained by solving an optimal control problem, which also guarantees that the parasitoid population remains above a desired level. Nevertheless, if there is a stable coexistence (i.e., an asymptotically stable positive equilibrium) and the equilibrium value of the parasitoid, x_2^* , is greater than the given threshold K, the desired level can be asymptotically reached without any further release. If the convergence is rapid enough, $x_2(t) > K$ can be reached very quickly, as shown in Figure 5 obtained with illustrative parameter values.

First, we find a necessary and sufficient condition for the existence of a positive equilibrium of system (1). To this end, we define function $f : R^3 \to R^3$,

$$f(x_1, x_2, x_3) = \begin{pmatrix} x_1(m_1 - \alpha \cdot x_1 - \gamma_1 \cdot x_3) \\ x_2(m_2 - \gamma_2 \cdot x_3) \\ x_3(-m_3 + \overline{\gamma}_1 \cdot x_1 + \overline{\gamma}_2 \cdot x_2) \end{pmatrix}.$$

Then, a nonzero vector $x^* \in R^3$ represents equilibrium for the considered dynamics if and only if $f(x^*) = 0$. The unique solution of this equation is easily obtained:

$$x_1^* = \frac{m_1\gamma_2 - m_2\gamma_1}{\alpha\gamma_2},$$
$$x_2^* = \frac{m_3\alpha\gamma_2 - \bar{\gamma}_1(m_1\gamma_2 - m_2\gamma_1)}{\alpha\gamma_2\bar{\gamma}_2},$$
$$x_3^* = \frac{m_2}{\gamma_2}.$$

Thus, a necessary and sufficient condition for $x^* > 0$ is

$$\frac{m_3 \alpha \gamma_2}{\overline{\gamma}_1} > m_1 \gamma_2 - m_2 \gamma_1 > 0 \tag{3}$$

To analyse the stability of system (1), we calculate the Jacobian

$$A = f'(x^*) = \begin{pmatrix} -\alpha x_1^* & 0 & -\gamma_1 x_1^* \\ 0 & 0 & -\gamma_2 x_2^* \\ \overline{\gamma_1} x_3^* & \overline{\gamma_2} x_3^* & 0 \end{pmatrix}$$

the characteristic polynomial of which is

$$\lambda^3 + \alpha x_1^* \lambda^2 + \left(\gamma_1 \overline{\gamma}_1 x_1^* + \gamma_2 \overline{\gamma}_2 x_2^* \right) x_3^* \lambda + \alpha \gamma_2 \overline{\gamma}_2 x_1^* x_2^* x_3^* = 0$$

Hence, by the Routh-Hurwitz criterion (see, e.g., Chen et al., 2004), for the asymptotic stability of equilibrium x^* it is enough to see that

$$\alpha x_1^* \Big(\gamma_1 \overline{\gamma}_1 x_1^* + \gamma_2 \overline{\gamma}_2 x_2^* \Big) x_3^* > \alpha \gamma_2 \overline{\gamma}_2 x_1^* x_2^* x_3^* ,$$

which always holds whenever equilibrium x^* is positive. We can conclude that condition (3) implies stable coexistence in the pest-parasitoid-predator system (1).

It is easy to confirm that for the parameter values obtained in Table 1, the sufficient conditions (3) for the stable coexistence are not satisfied. Nevertheless, the following example shows that the pest-parasitoid-predator system (1) with appropriate biological parameters display stable coexistence, and the parasitoid density even without control reaches the required density rather quickly.

Example. Consider the parameter system $m_1 = 1.54$; $m_2 = 1.73$; $m_3 = 0.1$; $\alpha = 0.23$; $\gamma_1 = 1.14$; $\gamma_2 = 1.39$; $\overline{\gamma}_1 = 0.16$ and $\overline{\gamma}_2 = 0.02$. A simple calculation shows that these parameters satisfy condition (3), therefore, system (1) has an asymptotically stable equilibrium $x^* = (0.527, 0.797, 1.245)$. The Jacobian

$$A := f'(x^*) = \begin{pmatrix} -0.12 & 0 & -0.6 \\ 0 & 0 & -1.09 \\ 0.2 & 0.02 & 0 \end{pmatrix}$$

has eigenvalues $\lambda_1 = -0.0492 + 0.3770i$, $\lambda_2 = -0.0492 - 0.3770i$, and $\lambda_3 = -0.0228$, all with negative real parts. Thus, the solutions tend towards a positive equilibrium, as demonstrated by the simulation results plotted in Figure 5. Setting K = 0.6, we can see that parasitoid density $x_2(t)$ exceeds the threshold K.

Figure 5: Solution of system (1) with initial value x(0) = (0.7, 0.4, 1.0), and exceeding the threshold value *K*=0.6 from time t_0 .



4. Discussion

Plants have developed different defence mechanisms against the phytophagous organisms that prey on them (e.g., Smith 2005). However, this also has an impact on subsequent trophic levels, such as entomophagous predators and parasitoids (e.g., Price et al. 1980). All of this is related to the different botanical species and their effects on both phytophagous and entomophagous organisms. However, few studies have investigated the differences among different cultivars from the same botanical species. Variations between cultivars can play an important role in the activity and efficiency of entomophagous agents (Bergman and Tingey 1979) as discussed by Bottrell et. al (1998), though questions remain, especially with respect to multitrophic effects involving complex interactions.

Our results demonstrating the effects of tomato cultivars on biological activity, especially on *N. tenuis* and, to a lesser extent, on *T. achaeae*, are supported by previous studies (e.g., Cabello et al. 2012, Vila et al. 2012). Thus, our results corroborate these findings.

Additionally, mathematical models similar to those shown here have been used to evaluate the efficacies of different species of *Trichogramma* in greenhouse tomatoes

(e.g., Gallego et al., 2013). This demonstrates the commercial usefulness of these mathematical models in the biological control of pest species.

< Notably, the studies carried out to elucidate the IGP in greenhouse tomato crops with a two-prey-one-predator model, specifically with the species *N. tenuis*, *T. achaeae* and *T. absoluta*, reveal the detrimental effects of the predator on the activity of the parasitoid and its resulting control of the pest prey *T. absoluta* (Gallego et al., 2013). The innovation of the present results is that they highlight that when the predator *N.tenuis* is used for the biological control on a pest species *B. tabaci*, different from the one than controls biologically *T.achaeae*, these detrimental effects are also produced.

Our results are unique because they modify existing models to study the varietal effects of both natural enemies. These effects can determine the efficiency of the generalist programmes of biological control at a practical level.

Further, this mathematical approach makes it possible to minimize the cost of applying agents, emphasizing the importance of the proposed optimal control model in integrated pest management (IPM) and/or biological control programmes.

5. Conclusion

The impact and efficiency of biological control programmes on pest species in greenhouse crops, in which various natural enemies are used during the crop cycle, can be influenced by the use of cultivars.

The optimal control model introduced here contributes to the methodological development of the biological control of pests in greenhouse crops.

This mathematical approach also makes it possible to minimize the cost of the application of agents, demonstrating the importance of the proposed optimal control model in integrated pest management (IPM).

Based on previous studies, this is the first example of a mathematical model based on an experiment that represents to this type of intraguild predation (IGP).

References

- Abrams P.A. (2012) Predator-prey models. In: Hastings, A. & Gross, L. (eds.). *Encyclopedia of theoretical ecology*. (pp. 587-594). Berkeley, USA: University of California Press.
- Banga, J.R., Balsa-Canto, E., Moles, C.G. & Alonso A.A. (2005). Dynamic Optimization of Bioprocesses: Efficient and Robust Numerical Strategies. Journal of Biothecnology 117, 407–419.
- Bardsley W.G. (2017) "SIMFIT statistical package", version 7.3.0. http://www.simfit.org.uk.
- Berryman A.A. (2004) The theoretical foundations of biological control. In Hawkins
 B.A. & Cornell H.V. (eds.) Theoretical approaches to biological control (pp.: 3-21). Cambridge, UK, Cambridge University Press.

- Cabello T., Bonfil F., Gallego J.R., Fernandez F.J., Gamez, M. & Garay J. (2015) Can interactions between and omnivorous hemipteran and an egg parasitoid limit the level of biological control for the tomato pinworm? Environ. Entomol. 44: 12-26.
- Cabello T., Gallego J.R., Vila E., Soler A., Pino del M., Carnero A., Hernandez-Suarez E. & Polaszek A. (2009) Biological control of the South American tomato pinworm, *Tuta absoluta* (Lep.: Gelechiidae), with releases of *Trichogramma achaeae* (Hym.: Trichogrammatidae) in tomato greenhouses of Spain. IOBC WPRS Bull. 49: 225–230.
- Cabello T., Gallego, J.R., Fernandez F.J., Gamez, M., Vila, E., Pino, M. & Hernandez, E. (2012). Biological control strategies for the South American tomato moth in greenhouse tomatoes. J. Econ. Entomol. 105: 2085-2096.
- Calvo F.J., Bolckmans K. & Belda J.E. (2012) Release rate for a pre-plant application of *Nesidiocoris tenuis* for *Bemisia tabaci* control in tomato. Biocontrol 57: 809-817.
- Castañe C., Alomar O., Goula M. & Gabarra R. (2004). Colonization of tomato greenhouses by the predatory mirid bugs *Macrolophus caliginosus* and *Dicyphus tamaninii*. Biol. Control 30: 591-597.
- Chen B.M., Lin, Z. & Shamesh Y. (2004) Linear Systems Theory. A Structural Decomposition Approach, Birkhauser, Boston.
- Gallego J.R., Lopez I., Gamez M., Cabello T., Varga Z. & Garay, J. (2013). Simulation model applied to biological pest control by entomophagous species in commercial tomato greenhouses. Hung. Agric. Eng. 25: 67-70.
- Hawkins, B. A. & Cornell, H. V. (eds) (2004). Theoretical approaches to biological control. Cambridge, UK: Cambridge Univ. Press.
- Hirmajer T., Balsa-Canto E. & Banga, J. (2009). DOTcvpSB, a Software Toolbox for Dynamic Optimization in Systems Biology. BMC Bioinformatics 10, 199.
- Holt R.D. and Polis G.A. (1997) A theoretical framework for intraguild predation. Am. Nat. 149: 745-764.
- IBM. 2014. IBM SPSS Statistics for Windows, Version 23.0 IBM Corp. Armonk, NY: IBM Corp.
- Janssen A., Pallini A., Venzon M & Sabelis M.W. (1998) Behaviour and indirect interactions in food webs of plant-inhabiting arthropods. Exp. Appl. Acarol. 22: 497-521.
- Messelink J.G., Bloemhard C.M.J., Sabelis M.W. & Janssen A. (2013) Biological control of aphids in the presence of thrips and their enemies. Biocontrol 58: 45-55.
- Oliveira L., Durao A.C., Fontes J., Roja I.S. & Tavares J. (2017) Potential of *Trichogramma achaeae* (Hym.: Trichogrammatidae) in biological bontrol of *Tuta absoluta* (Lep.: Gelechiidae) in Azorean greenhouse tomato crops. J. Econ. Entomol. 110: 2010-2015.
- Polaszek, A., Rugman-Jones P.F., Stouthamer R., Hernandez-Suarez E., Cabello T. & Pino M. del (2012) Molecular and morphological diagnoses of Canary Islands

Trichogramma species for biocontrol of *Chrysodeixis chalcites* (Lep. Noctuidae) and *Tuta absoluta* (Lep. Gelechiidae). BioControl 57: 21–35.

- Price P.W., Bouton C.E., Gross P., McPheron B.A., Thompson J.N. & Weis A.E. (1980). Influence of plant on interactions between insect herbivores and natural enemies. Ann. Rev. Ecol. Syst. 11: 41-65.
- Rosenheim J.A. & Harmon J.P. (2006) The influence of intraguild predation on the suppression of a shared prey population: an empirical reassessment, pp: 1-20. In Brodeur J. & Boivin G. (eds.), Trophic and guild interactions in biological control. Springer,Dordrecht, NL.
- Ruppel R.F. (1983) Cumulative insect-days as an index of crop protection. J. Econ. Entomol. 76: 375-377.
- Sanchez, J. A. 2008. Zoophytophagy in the plantbug *Nesidiocoris tenuis*. Agric. For. Entomol. 10: 75–80.
- Smith C.M. (2005) Plant resistance to arthropods: molecular and conventional approaches. Dordrecht, NL: Springer.
- Varga Z., Sebestyen Z., Gamez M., Cabello T. & Attias A. (2010), Models of applied population dynamics. Mech. Eng. Lett. 10: 22-36.
- Vila, E. & Cabello, T. (2014). Biosystems engineering applied to greenhouse pest control. In Torres, I., Guevara, R. (eds.) Biosystems engineering: Biofactories for food production in the XXI Century (pp.: 99-128). Germany, Springer.
- Vila, E., Parra, A., Beltran, D., Gallego, J.R., Fernandez, F.J. & Cabello, T. (2012). IPM strategies in tomato crops in Spanish greenhouses: Effects of cultivars and the integration of natural enemies. IOBC-WPRS Bull. 80: 245-251.
- Yano E. (2005) Effects of intraguild predation and interspecific competition among biological control agents in augmentative biological control in greenhouses, pp. 523-530. In Hoddle M.S. (ed.), Second International Symposium on Biological Control of Arthropods, September 12-16, Davos, CH. USDA Forest Service (Publication FHTET-2005-08).